

Chapter 13

Eigenvectors and Eigenvalues

13.1 Eigenvectors and Eigenvalues of a Linear Map

Given a finite-dimensional vector space E , let $f: E \rightarrow E$ be any linear map. If, by luck, there is a basis (e_1, \dots, e_n) of E with respect to which f is represented by a *diagonal matrix*

$$D = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & \vdots \\ \vdots & \dots & \dots & 0 \\ 0 & \dots & 0 & \lambda_n \end{pmatrix},$$

then the action of f on E is very simple; in every “direction” e_i , we have

$$f(e_i) = \lambda_i e_i.$$

We can think of f as a transformation that *stretches or shrinks* space along the direction e_1, \dots, e_n (at least if E is a real vector space).

In terms of matrices, the above property translates into the fact that there is an invertible matrix P and a diagonal matrix D such that a matrix A can be factored as

$$A = PDP^{-1}.$$

When this happens, we say that f (or A) is *diagonalizable*, the λ_i s are called the *eigenvalues* of f , and the e_i s are *eigenvectors* of f .

For example, we will see that *every symmetric matrix can be diagonalized*.

Unfortunately, *not every matrix can be diagonalized*.

For example, the matrix

$$A_1 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

can't be diagonalized.

Sometimes, a matrix fails to be diagonalizable because its eigenvalues do not belong to the field of coefficients, such as

$$A_2 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

whose eigenvalues are $\pm i$.

This is not a serious problem because A_2 can be diagonalized over the complex numbers.

However, A_1 is a “fatal” case! Indeed, its eigenvalues are both 1 and the problem is that A_1 *does not have enough eigenvectors to span E* .

The next best thing is that there is a basis with respect to which f is represented by an *upper triangular* matrix.

In this case we say that f can be *triangularized*.

As we will see in Section 13.2, if all the eigenvalues of f belong to the field of coefficients K , then f can be triangularized. In particular, this is the case if $K = \mathbb{C}$.

Now, an alternative to triangularization is to consider the representation of f with respect to *two* bases (e_1, \dots, e_n) and (f_1, \dots, f_n) , rather than a single basis.

In this case, if $K = \mathbb{R}$ or $K = \mathbb{C}$, it turns out that we can even pick these bases to be *orthonormal*, and we get a diagonal matrix Σ with *nonnegative entries*, such that

$$f(e_i) = \sigma_i f_i, \quad 1 \leq i \leq n.$$

The nonzero σ_i s are the *singular values* of f , and the corresponding representation is the *singular value decomposition*, or *SVD*.

The notion of eigenvalue of a linear map $f: E \rightarrow E$ defined on an infinite-dimensional space E is quite subtle because it cannot be defined in terms of eigenvectors as in the finite-dimensional case.

The problem is that the map $\lambda \text{id} - f$ (with $\lambda \in \mathbb{C}$) could be noninvertible (because it is not surjective) and yet injective.

In finite dimension this cannot happen, so until further notice we *assume that E is of finite dimension n .*

Definition 13.1. Given any vector space E of finite dimension n and any linear map $f: E \rightarrow E$, a scalar $\lambda \in K$ is called an *eigenvalue, or proper value, or characteristic value of f* if there is some *nonzero* vector $u \in E$ such that

$$f(u) = \lambda u.$$

Equivalently, λ is an eigenvalue of f if $\text{Ker}(\lambda \text{id} - f)$ is nontrivial (i.e., $\text{Ker}(\lambda \text{id} - f) \neq \{0\}$).

A vector $u \in E$ is called an *eigenvector, or proper vector, or characteristic vector of f* if $u \neq 0$ and if there is some $\lambda \in K$ such that

$$f(u) = \lambda u;$$

the scalar λ is then an eigenvalue, and we say that u is an *eigenvector associated with λ* .

Given any eigenvalue $\lambda \in K$, the nontrivial subspace $\text{Ker}(\lambda \text{id} - f)$ consists of all the eigenvectors associated with λ together with the zero vector; this subspace is denoted by $E_\lambda(f)$, or $E(\lambda, f)$, or even by E_λ , and is called the *eigenspace associated with λ , or proper subspace associated with λ* .

Remark: As we emphasized in the remark following Definition 7.4, we *require an eigenvector to be nonzero*.

This requirement seems to have more benefits than inconvenients, even though it may be considered somewhat inelegant because the set of all eigenvectors associated with an eigenvalue is not a subspace since the zero vector is excluded.

Note that distinct eigenvectors may correspond to the same eigenvalue, but distinct eigenvalues correspond to disjoint sets of eigenvectors.

Proposition 13.1. *Let E be any vector space of finite dimension n and let f be any linear map $f: E \rightarrow E$. The eigenvalues of f are the roots (in K) of the polynomial*

$$\det(\lambda \text{id} - f).$$

Proof. A scalar $\lambda \in K$ is an eigenvalue of f iff there is some vector $u \neq 0$ in E such that

$$f(u) = \lambda u$$

iff

$$(\lambda \text{id} - f)(u) = 0$$

iff $(\lambda \text{id} - f)$ is not invertible

iff by Proposition 5.15,

$$\det(\lambda \text{id} - f) = 0. \quad \square$$

Definition 13.2. Given any vector space E of dimension n , for any linear map $f: E \rightarrow E$, the polynomial $P_f(X) = \chi_f(X) = \det(X \text{id} - f)$ is called the *characteristic polynomial of f* . For any square matrix A , the polynomial $P_A(X) = \chi_A(X) = \det(XI - A)$ is called the *characteristic polynomial of A* .

Note that we already encountered the characteristic polynomial in Section 5.7; see Definition 5.11.

Given any basis (e_1, \dots, e_n) , if $A = M(f)$ is the matrix of f w.r.t. (e_1, \dots, e_n) , we can compute the characteristic polynomial $\chi_f(X) = \det(X \text{id} - f)$ of f by expanding the following determinant:

$$\det(XI - A) = \begin{vmatrix} X - a_{11} & -a_{12} & \cdots & -a_{1n} \\ -a_{21} & X - a_{22} & \cdots & -a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ -a_{n1} & -a_{n2} & \cdots & X - a_{nn} \end{vmatrix}.$$

If we expand this determinant, we find that

$$\begin{aligned} \chi_A(X) = \det(XI - A) &= X^n - (a_{11} + \cdots + a_{nn})X^{n-1} \\ &\quad + \cdots + (-1)^n \det(A). \end{aligned}$$

The sum $\text{tr}(A) = a_{11} + \cdots + a_{nn}$ of the diagonal elements of A is called the *trace of A* .

Since the characteristic polynomial depends only on f , $\text{tr}(A)$ has the same value for all matrices A representing f . We let $\text{tr}(f) = \text{tr}(A)$ be the *trace* of f .

Remark: The characteristic polynomial of a linear map is sometimes defined as $\det(f - X \text{id})$. Since

$$\det(f - X \text{id}) = (-1)^n \det(X \text{id} - f),$$

this makes essentially no difference but the version $\det(XI - f)$ has the small advantage that the coefficient of X^n is $+1$.

If we write

$$\begin{aligned} \chi_A(X) = \det(XI - A) &= X^n - \tau_1(A)X^{n-1} \\ &+ \cdots + (-1)^k \tau_k(A)X^{n-k} + \cdots + (-1)^n \tau_n(A), \end{aligned}$$

then we just proved that

$$\tau_1(A) = \text{tr}(A) \quad \text{and} \quad \tau_n(A) = \det(A).$$

It is also possible to express $\tau_k(A)$ in terms of determinants of certain submatrices of A .

For any nonempty ordered subset, $I \subseteq \{1, \dots, n\}$, say $I = \{i_1 < \dots < i_k\}$, let $A_{I,I}$ be the $k \times k$ submatrix of A obtained by first selecting the columns whose indices belong to I , and then the rows whose indices also belong to I .

Then, it can be shown that

$$\tau_k(A) = \sum_{\substack{I \subseteq \{1, \dots, n\} \\ I = \{i_1, \dots, i_k\} \\ i_1 < \dots < i_k}} \det(A_{I,I}).$$

If all the roots, $\lambda_1, \dots, \lambda_n$, of the polynomial $\det(XI - A)$ belong to the field K , then we can write

$$\det(XI - A) = (X - \lambda_1) \cdots (X - \lambda_n),$$

where some of the λ_i s may appear more than once.

Consequently,

$$\begin{aligned}\chi_A(X) = \det(XI - A) &= X^n - \sigma_1(\lambda)X^{n-1} \\ &+ \cdots + (-1)^k \sigma_k(\lambda)X^{n-k} + \cdots + (-1)^n \sigma_n(\lambda),\end{aligned}$$

where

$$\sigma_k(\lambda) = \sum_{\substack{I \subseteq \{1, \dots, n\} \\ |I|=k}} \prod_{i \in I} \lambda_i,$$

the *k*th elementary symmetric polynomial (or function) of the λ_i 's, with $\lambda = (\lambda_1, \dots, \lambda_n)$.

For $n = 5$, the elementary symmetric polynomials are listed below:

$$\sigma_0(\lambda) = 1$$

$$\sigma_1(\lambda) = \lambda_1 + \lambda_2 + \lambda_3 + \lambda_4 + \lambda_5$$

$$\begin{aligned} \sigma_2(\lambda) = & \lambda_1\lambda_2 + \lambda_1\lambda_3 + \lambda_1\lambda_4 + \lambda_1\lambda_5 + \lambda_2\lambda_3 + \lambda_2\lambda_4 + \lambda_2\lambda_5 \\ & + \lambda_3\lambda_4 + \lambda_3\lambda_5 + \lambda_4\lambda_5 \end{aligned}$$

$$\begin{aligned} \sigma_3(\lambda) = & \lambda_3\lambda_4\lambda_5 + \lambda_2\lambda_4\lambda_5 + \lambda_2\lambda_3\lambda_5 + \lambda_2\lambda_3\lambda_4 + \lambda_1\lambda_4\lambda_5 \\ & + \lambda_1\lambda_3\lambda_5 + \lambda_1\lambda_3\lambda_4 + \lambda_1\lambda_2\lambda_5 + \lambda_1\lambda_2\lambda_4 + \lambda_1\lambda_2\lambda_3 \end{aligned}$$

$$\begin{aligned} \sigma_4(\lambda) = & \lambda_1\lambda_2\lambda_3\lambda_4 + \lambda_1\lambda_2\lambda_3\lambda_5 + \lambda_1\lambda_2\lambda_4\lambda_5 \\ & + \lambda_1\lambda_3\lambda_4\lambda_5 + \lambda_2\lambda_3\lambda_4\lambda_5 \end{aligned}$$

$$\sigma_5(\lambda) = \lambda_1\lambda_2\lambda_3\lambda_4\lambda_5.$$

Since

$$\begin{aligned} \chi_A(X) &= X^n - \tau_1(A)X^{n-1} + \cdots + (-1)^k \tau_k(A)X^{n-k} \\ &\quad + \cdots + (-1)^n \tau_n(A) \\ &= X^n - \sigma_1(\lambda)X^{n-1} + \cdots + (-1)^k \sigma_k(\lambda)X^{n-k} \\ &\quad + \cdots + (-1)^n \sigma_n(\lambda), \end{aligned}$$

we have

$$\sigma_k(\lambda) = \tau_k(A), \quad k = 1, \dots, n.$$

In particular, the product of the eigenvalues of f is equal to $\det(A) = \det(f)$, and the sum of the eigenvalues of f is equal to the trace $\operatorname{tr}(A) = \operatorname{tr}(f)$, of f .

For the record,

$$\begin{aligned}\operatorname{tr}(f) &= \lambda_1 + \cdots + \lambda_n \\ \det(f) &= \lambda_1 \cdots \lambda_n,\end{aligned}$$

where $\lambda_1, \dots, \lambda_n$ are the eigenvalues of f (and A), where some of the λ_i s may appear more than once.

In particular, f is not invertible iff it admits 0 as an eigenvalue.

Remark: Depending on the field K , the characteristic polynomial $\chi_A(X) = \det(XI - A)$ may or may not have roots in K .

This motivates considering *algebraically closed fields*. For example, over $K = \mathbb{R}$, not every polynomial has real roots. For example, for the matrix

$$A = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix},$$

the characteristic polynomial $\det(XI - A)$ has no real roots unless $\theta = k\pi$.

However, over the field \mathbb{C} of complex numbers, every polynomial has roots. For example, the matrix above has the roots $\cos \theta \pm i \sin \theta = e^{\pm i\theta}$.

Definition 13.3. Let A be an $n \times n$ matrix over a field, K . Assume that all the roots of the characteristic polynomial $\chi_A(X) = \det(XI - A)$ of A belong to K , which means that we can write

$$\det(XI - A) = (X - \lambda_1)^{k_1} \cdots (X - \lambda_m)^{k_m},$$

where $\lambda_1, \dots, \lambda_m \in K$ are the distinct roots of $\det(XI - A)$ and $k_1 + \cdots + k_m = n$.

The integer, k_i , is called the *algebraic multiplicity* of the eigenvalue λ_i and the dimension of the eigenspace, $E_{\lambda_i} = \text{Ker}(\lambda_i I - A)$, is called the *geometric multiplicity* of λ_i . We denote the algebraic multiplicity of λ_i by $\text{alg}(\lambda_i)$ and its geometric multiplicity by $\text{geo}(\lambda_i)$.

By definition, the sum of the algebraic multiplicities is equal to n but the sum of the geometric multiplicities can be strictly smaller.

Proposition 13.2. *Let A be an $n \times n$ matrix over a field K and assume that all the roots of the characteristic polynomial $\chi_A(X) = \det(XI - A)$ of A belong to K . For every eigenvalue λ_i of A , the geometric multiplicity of λ_i is always less than or equal to its algebraic multiplicity, that is,*

$$\text{geo}(\lambda_i) \leq \text{alg}(\lambda_i).$$

Proposition 13.3. *Let E be any vector space of finite dimension n and let f be any linear map. If u_1, \dots, u_m are eigenvectors associated with pairwise distinct eigenvalues $\lambda_1, \dots, \lambda_m$, then the family (u_1, \dots, u_m) is linearly independent.*

Thus, from Proposition 13.3, if $\lambda_1, \dots, \lambda_m$ are all the pairwise distinct eigenvalues of f (where $m \leq n$), we have a *direct sum*

$$E_{\lambda_1} \oplus \cdots \oplus E_{\lambda_m}$$

of the eigenspaces E_{λ_i} .

Unfortunately, it is not always the case that

$$E = E_{\lambda_1} \oplus \cdots \oplus E_{\lambda_m}.$$

When

$$E = E_{\lambda_1} \oplus \cdots \oplus E_{\lambda_m},$$

we say that f is *diagonalizable* (and similarly for any matrix associated with f).

Indeed, picking a basis in each E_{λ_i} , we obtain a matrix which is a diagonal matrix consisting of the eigenvalues, each λ_i occurring a number of times equal to the dimension of E_{λ_i} .

This happens if the algebraic multiplicity and the geometric multiplicity of every eigenvalue are equal.

In particular, when the characteristic polynomial has n *distinct roots*, then f is diagonalizable.

It can also be shown that symmetric matrices have real eigenvalues and can be diagonalized.

For a negative example, we leave as exercise to show that the matrix

$$M = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

cannot be diagonalized, even though 1 is an eigenvalue.

The problem is that the eigenspace of 1 only has dimension 1.

The matrix

$$A = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

cannot be diagonalized either, because it has no real eigenvalues, unless $\theta = k\pi$.

However, over the field of complex numbers, it can be diagonalized.

13.2 Reduction to Upper Triangular Form

Unfortunately, not every linear map on a complex vector space can be diagonalized.

The next best thing is to “triangularize,” which means to find a basis over which the matrix has zero entries below the main diagonal.

Fortunately, such a basis always exist.

We say that a square matrix A is an *upper triangular matrix* if it has the following shape,

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n-1} & a_{1n} \\ 0 & a_{22} & a_{23} & \cdots & a_{2n-1} & a_{2n} \\ 0 & 0 & a_{33} & \cdots & a_{3n-1} & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & a_{n-1n-1} & a_{n-1n} \\ 0 & 0 & 0 & \cdots & 0 & a_{nn} \end{pmatrix},$$

i.e., $a_{ij} = 0$ whenever $j < i$, $1 \leq i, j \leq n$.

Theorem 13.4. *Given any finite dimensional vector space over a field K , for any linear map $f: E \rightarrow E$, there is a basis (u_1, \dots, u_n) with respect to which f is represented by an upper triangular matrix (in $M_n(K)$) iff all the eigenvalues of f belong to K . Equivalently, for every $n \times n$ matrix $A \in M_n(K)$, there is an invertible matrix P and an upper triangular matrix T (both in $M_n(K)$) such that*

$$A = PTP^{-1}$$

iff all the eigenvalues of A belong to K .

If $A = PTP^{-1}$ where T is upper triangular, note that the diagonal entries of T are the eigenvalues $\lambda_1, \dots, \lambda_n$ of A .

Also, if A is a real matrix whose eigenvalues are all real, then P can be chosen to real, and if A is a rational matrix whose eigenvalues are all rational, then P can be chosen rational.

Since any polynomial over \mathbb{C} has all its roots in \mathbb{C} , Theorem 13.4 implies that *every complex $n \times n$ matrix can be triangularized*.

If λ is an eigenvalue of the matrix A and if u is an eigenvector associated with λ , from

$$Au = \lambda u,$$

we obtain

$$A^2u = A(Au) = A(\lambda u) = \lambda Au = \lambda^2u,$$

which shows that λ^2 is an eigenvalue of A^2 for the eigenvector u .

An obvious induction shows that λ^k is an eigenvalue of A^k for the eigenvector u , for all $k \geq 1$.

Now, if all eigenvalues $\lambda_1, \dots, \lambda_n$ of A are in K , it follows that $\lambda_1^k, \dots, \lambda_n^k$ are eigenvalues of A^k .

However, it is not obvious that A^k does not have other eigenvalues. In fact, this can't happen, and this can be proved using Theorem 13.4.

Proposition 13.5. *Given any $n \times n$ matrix $A \in M_n(K)$ with coefficients in a field K , if all eigenvalues $\lambda_1, \dots, \lambda_n$ of A are in K , then for every polynomial $q(X) \in K[X]$, the eigenvalues of $q(A)$ are exactly $(q(\lambda_1), \dots, q(\lambda_n))$.*

If E is a Hermitian space, the proof of Theorem 13.4 can be easily adapted to prove that there is an *orthonormal* basis (u_1, \dots, u_n) with respect to which the matrix of f is upper triangular. This is usually known as *Schur's lemma*.

Theorem 13.6. (*Schur decomposition*) *Given any linear map $f: E \rightarrow E$ over a complex Hermitian space E , there is an orthonormal basis (u_1, \dots, u_n) with respect to which f is represented by an upper triangular matrix. Equivalently, for every $n \times n$ matrix $A \in M_n(\mathbb{C})$, there is a unitary matrix U and an upper triangular matrix T such that*

$$A = UTU^*.$$

If A is real and if all its eigenvalues are real, then there is an orthogonal matrix Q and a real upper triangular matrix T such that

$$A = QTQ^T.$$

Using, Theorem 13.6, we can derive two very important results.

Proposition 13.7. *If A is a Hermitian matrix (i.e. $A^* = A$), then its eigenvalues are real and A can be diagonalized with respect to an orthonormal basis of eigenvectors. In matrix terms, there is a unitary matrix U and a real diagonal matrix D such that $A = UDU^*$. If A is a real symmetric matrix (i.e. $A^\top = A$), then its eigenvalues are real and A can be diagonalized with respect to an orthonormal basis of eigenvectors. In matrix terms, there is an orthogonal matrix Q and a real diagonal matrix D such that $A = QDQ^\top$.*

When a real matrix A has complex eigenvalues, there is a version of Theorem 13.6 involving only real matrices provided that we allow T to be block upper-triangular (the diagonal entries may be 2×2 matrices or real entries).

Theorem 13.6 is not a very practical result but it is a useful theoretical result to cope with matrices that cannot be diagonalized.

For example, it can be used to prove that *every complex matrix is the limit of a sequence of diagonalizable matrices that have distinct eigenvalues!*

13.3 Location of Eigenvalues

If A is an $n \times n$ complex (or real) matrix A , it would be useful to know, even roughly, where the eigenvalues of A are located in the complex plane \mathbb{C} .

The Gershgorin discs provide some precise information about this.

Definition 13.4. For any complex $n \times n$ matrix A , for $i = 1, \dots, n$, let

$$R'_i(A) = \sum_{\substack{j=1 \\ j \neq i}}^n |a_{ij}|$$

and let

$$G(A) = \bigcup_{i=1}^n \{z \in \mathbb{C} \mid |z - a_{ii}| \leq R'_i(A)\}.$$

Each disc $\{z \in \mathbb{C} \mid |z - a_{ii}| \leq R'_i(A)\}$ is called a *Gershgorin disc* and their union $G(A)$ is called the *Gershgorin domain*.

Theorem 13.8. (*Gershgorin's disc theorem*) For any complex $n \times n$ matrix A , all the eigenvalues of A belong to the Gershgorin domain $G(A)$. Furthermore the following properties hold:

(1) If A is strictly row diagonally dominant, that is

$$|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}|, \quad \text{for } i = 1, \dots, n,$$

then A is invertible.

(2) If A is strictly row diagonally dominant, and if $a_{ii} > 0$ for $i = 1, \dots, n$, then every eigenvalue of A has a strictly positive real part.

In particular, Theorem 13.8 implies that if a symmetric matrix is strictly row diagonally dominant and has strictly positive diagonal entries, then it is positive definite.

Theorem 13.8 is sometimes called the *Gershgorin–Hadamard theorem*.

Since A and A^\top have the same eigenvalues (even for complex matrices) we also have a version of Theorem 13.8 for the discs of radius

$$C'_j(A) = \sum_{\substack{i=1 \\ i \neq j}}^n |a_{ij}|,$$

whose domain is denoted by $G(A^\top)$.

Theorem 13.9. *For any complex $n \times n$ matrix A , all the eigenvalues of A belong to the intersection of the Gershgorin domains, $G(A) \cap G(A^\top)$. Furthermore the following properties hold:*

(1) *If A is strictly column diagonally dominant, that is*

$$|a_{ii}| > \sum_{i=1, i \neq j}^n |a_{ij}|, \quad \text{for } j = 1, \dots, n,$$

then A is invertible.

(2) *If A is strictly column diagonally dominant, and if $a_{ii} > 0$ for $i = 1, \dots, n$, then every eigenvalue of A has a strictly positive real part.*

There are refinements of Gershgorin's theorem and eigenvalue location results involving other domains besides discs; for more on this subject, see Horn and Johnson [19], Sections 6.1 and 6.2.

Remark: Neither strict row diagonal dominance nor strict column diagonal dominance are necessary for invertibility. Also, if we relax all strict inequalities to inequalities, then row diagonal dominance (or column diagonal dominance) is not a sufficient condition for invertibility.

13.4 Conditioning of Eigenvalue Problems

The following $n \times n$ matrix

$$A = \begin{pmatrix} 0 & & & & & \\ 1 & 0 & & & & \\ & 1 & 0 & & & \\ & & \cdots & \cdots & & \\ & & & 1 & 0 & \\ & & & & 1 & 0 \end{pmatrix}$$

has the eigenvalue 0 with multiplicity n .

However, if we perturb the top rightmost entry of A by ϵ , it is easy to see that the characteristic polynomial of the matrix

$$A(\epsilon) = \begin{pmatrix} 0 & & & & \epsilon \\ 1 & 0 & & & \\ & 1 & 0 & & \\ & & \cdots & \cdots & \\ & & & 1 & 0 \\ & & & & 1 & 0 \end{pmatrix}$$

is $X^n - \epsilon$.

It follows that if $n = 40$ and $\epsilon = 10^{-40}$, $A(10^{-40})$ has the eigenvalues $e^{k2\pi i/40}10^{-1}$ with $k = 1, \dots, 40$.

Thus, we see that a very small change ($\epsilon = 10^{-40}$) to the matrix A causes a significant change to the eigenvalues of A (from 0 to $e^{k2\pi i/40}10^{-1}$).

Indeed, the relative error is 10^{-39} .

Worse, due to machine precision, since very small numbers are treated as 0, the error on the computation of eigenvalues (for example, of the matrix $A(10^{-40})$) can be very large.

This phenomenon is similar to the phenomenon discussed in Section 7.3 where we studied the effect of a small perturbation of the coefficients of a linear system $Ax = b$ on its solution.

In Section 7.3, we saw that the behavior of a linear system under small perturbations is governed by the condition number $\text{cond}(A)$ of the matrix A .

In the case of the eigenvalue problem (finding the eigenvalues of a matrix), we will see that the conditioning of the problem depends on the condition number of the change of basis matrix P used in reducing the matrix A to its diagonal form $D = P^{-1}AP$, rather than on the condition number of A itself.

The following proposition in which we assume that A is diagonalizable and that the matrix norm $\| \cdot \|$ satisfies a special condition (satisfied by the operator norms $\| \cdot \|_p$ for $p = 1, 2, \infty$), is due to Bauer and Fike (1960).

Proposition 13.10. *Let $A \in M_n(\mathbb{C})$ be a diagonalizable matrix, P be an invertible matrix and, D be a diagonal matrix $D = \text{diag}(\lambda_1, \dots, \lambda_n)$ such that*

$$A = PDP^{-1},$$

and let $\| \cdot \|$ be a matrix norm such that

$$\|\text{diag}(\alpha_1, \dots, \alpha_n)\| = \max_{1 \leq i \leq n} |\alpha_i|,$$

for every diagonal matrix. Then, for every perturbation matrix δA , if we write

$$B_i = \{z \in \mathbb{C} \mid |z - \lambda_i| \leq \text{cond}(P) \|\delta A\|\},$$

for every eigenvalue λ of $A + \delta A$, we have

$$\lambda \in \bigcup_{k=1}^n B_k.$$

Proposition 13.10 implies that for any diagonalizable matrix A , if we define $\Gamma(A)$ by

$$\Gamma(A) = \inf\{\text{cond}(P) \mid P^{-1}AP = D\},$$

then for every eigenvalue λ of $A + \delta A$, we have

$$\lambda \in \bigcup_{k=1}^n \{z \in \mathbb{C}^n \mid |z - \lambda_k| \leq \Gamma(A) \|\delta A\|\}.$$

Definition 13.5. The number $\Gamma(A)$ is called the *conditioning of A relative to the eigenvalue problem*.

If A is a normal matrix, since by Theorem 14.22, A can be diagonalized with respect to a unitary matrix U , and since for the spectral norm $\|U\|_2 = 1$, we see that $\Gamma(A) = 1$.

Therefore, normal matrices are very well conditioned w.r.t. the eigenvalue problem. In fact, for every eigenvalue λ of $A + \delta A$ (with A normal), we have

$$\lambda \in \bigcup_{k=1}^n \{z \in \mathbb{C}^n \mid |z - \lambda_k| \leq \|\delta A\|_2\}.$$

If A and $A + \delta A$ are both symmetric (or Hermitian), there are sharper results; see Proposition 14.28.

Note that the matrix $A(\epsilon)$ from the beginning of the section is not normal.

13.5 Eigenvalues of the Matrix Exponential

The Schur decomposition yields a characterization of the eigenvalues of the matrix exponential e^A in terms of the eigenvalues of the matrix A .

Proposition 13.11. *Let A and U be (real or complex) matrices and assume that U is invertible. Then*

$$e^{UAU^{-1}} = Ue^AU^{-1}.$$

Proposition 13.12. *Given any complex $n \times n$ matrix A , if $\lambda_1, \dots, \lambda_n$ are the eigenvalues of A , then $e^{\lambda_1}, \dots, e^{\lambda_n}$ are the eigenvalues of e^A . Furthermore, if u is an eigenvector of A for λ_i , then u is an eigenvector of e^A for e^{λ_i} .*

As a consequence, we obtain the following result.

Proposition 13.13. *For every complex (or real) square matrix A , we have*

$$\det(e^A) = e^{\operatorname{tr}(A)},$$

where $\operatorname{tr}(A)$ is the trace of A , i.e., the sum $a_{11} + \cdots + a_{nn}$ of its diagonal entries.

If B is a skew symmetric matrix, since $\operatorname{tr}(B) = 0$, we deduce that $\det(e^B) = e^0 = 1$. This allows us to obtain the following result. Recall that the (real) vector space of skew symmetric matrices is denoted by $\mathfrak{so}(n)$.

Proposition 13.14. *For every skew symmetric matrix $B \in \mathfrak{so}(n)$, we have $e^B \in \mathbf{SO}(n)$, that is, e^B is a rotation.*

Proposition 13.14 shows that the map $B \mapsto e^B$ is a map $\exp: \mathfrak{so}(n) \rightarrow \mathbf{SO}(n)$. It is not injective, but it can be shown (using one of the spectral theorems) that it is *surjective*.

If B is a (real) symmetric matrix, then

$$(e^B)^\top = e^{B^\top} = e^B,$$

so e^B is also symmetric.

Since the eigenvalues $\lambda_1, \dots, \lambda_n$ of B are real, by Proposition 13.12, since the eigenvalues of e^B are $e^{\lambda_1}, \dots, e^{\lambda_n}$ and the λ_i are real, we have $e^{\lambda_i} > 0$ for $i = 1, \dots, n$, which implies that e^B is *symmetric positive definite*.

In fact, it can be shown that for every symmetric positive definite matrix A , there is a *unique* symmetric matrix B such that $A = e^B$; see Gallier [14].

