

Fundamentals of Linear Algebra and Optimization

Jean Gallier

Homework 1

September 3 2013; Due September 17, 2013

Problem B1 (10 pts). Prove that the axioms of vector spaces imply that

$$\begin{aligned}\alpha \cdot 0 &= 0 \\ 0 \cdot v &= 0 \\ \alpha \cdot (-v) &= -(\alpha \cdot v) \\ (-\alpha) \cdot v &= -(\alpha \cdot v),\end{aligned}$$

for all $v \in E$ and all $\alpha \in K$, where E is a vector space over K .

Problem B2 (40 pts). (In solving this problem, **do not use determinants**). (1) Let (u_1, \dots, u_m) and (v_1, \dots, v_m) be two families of vectors in some vector space E . Assume that each v_i is a linear combination of the u_j s, so that

$$v_i = a_{i1}u_1 + \dots + a_{im}u_m, \quad 1 \leq i \leq m,$$

and that the matrix $A = (a_{ij})$ is an upper-triangular matrix, which means that if $1 \leq j < i \leq m$, then $a_{ij} = 0$. Prove that if (u_1, \dots, u_m) are linearly independent and if all the diagonal entries of A are nonzero, then (v_1, \dots, v_m) are also linearly independent.

Hint. Use induction on m .

(2) Let $A = (a_{ij})$ be an upper-triangular matrix. Prove that if all the diagonal entries of A are nonzero, then A is invertible and the inverse A^{-1} of A is also upper-triangular.

Hint. Use induction on m .

Prove that if A is invertible, then all the diagonal entries of A are nonzero (do not use determinants or eigenvalues!).

(3) Prove that if the families (u_1, \dots, u_m) and (v_1, \dots, v_m) are related as in (1), then (u_1, \dots, u_m) are linearly independent iff (v_1, \dots, v_m) are.

Problem B3 (40 pts). (In solving this problem, **do not use determinants**). Consider

the $n \times n$ matrix

$$A = \begin{pmatrix} 1 & 2 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 2 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & 2 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 0 & 1 & 2 & 0 \\ 0 & 0 & \dots & 0 & 0 & 1 & 2 \\ 0 & 0 & \dots & 0 & 0 & 0 & 1 \end{pmatrix}.$$

(1) Find the solution $x = (x_1, \dots, x_n)$ of the linear system

$$Ax = b,$$

for

$$b = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{pmatrix}.$$

(2) Prove that the matrix A is invertible and find its inverse A^{-1} . Given that the number of atoms in the universe is estimated to be $\leq 10^{82}$, explain why it is practically impossible that the inverse of A will ever be computed if $n \geq 300$.

(3) Assume b is perturbed by a small amount δb (note that δb is a vector). Find the new solution of the system

$$A(x + \delta x) = b + \delta b,$$

where δx is also a vector. In the case where $b = (0, \dots, 0, 1)$, and $\delta b = (0, \dots, 0, \epsilon)$, show that

$$|(\delta x)_1| = 2^{n-1}|\epsilon|.$$

(where $|(\delta x)_1|$ is the first component of δx).

(4) Prove that $(A - I)^n = 0$.

Problem B4 (80 pts). Consider the polynomials

$$\begin{aligned} B_0^2(t) &= (1-t)^2 & B_1^2(t) &= 2(1-t)t & B_2^2(t) &= t^2 \\ B_0^3(t) &= (1-t)^3 & B_1^3(t) &= 3(1-t)^2t & B_2^3(t) &= 3(1-t)t^2 & B_3^3(t) &= t^3, \end{aligned}$$

known as the *Bernstein polynomials* of degree 2 and 3.

(1) Show that the Bernstein polynomials $B_0^2(t), B_1^2(t), B_2^2(t)$ are expressed as linear combinations of the basis $(1, t, t^2)$ of the vector space of polynomials of degree at most 2 as follows:

$$\begin{pmatrix} B_0^2(t) \\ B_1^2(t) \\ B_2^2(t) \end{pmatrix} = \begin{pmatrix} 1 & -2 & 1 \\ 0 & 2 & -2 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ t \\ t^2 \end{pmatrix}.$$

Prove that

$$B_0^2(t) + B_1^2(t) + B_2^2(t) = 1.$$

(2) Show that the Bernstein polynomials $B_0^3(t), B_1^3(t), B_2^3(t), B_3^3(t)$ are expressed as linear combinations of the basis $(1, t, t^2, t^3)$ of the vector space of polynomials of degree at most 3 as follows:

$$\begin{pmatrix} B_0^3(t) \\ B_1^3(t) \\ B_2^3(t) \\ B_3^3(t) \end{pmatrix} = \begin{pmatrix} 1 & -3 & 3 & -1 \\ 0 & 3 & -6 & 3 \\ 0 & 0 & 3 & -3 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ t \\ t^2 \\ t^3 \end{pmatrix}.$$

Prove that

$$B_0^3(t) + B_1^3(t) + B_2^3(t) + B_3^3(t) = 1.$$

(3) Prove that the Bernstein polynomials of degree 2 are linearly independent, and that the Bernstein polynomials of degree 3 are linearly independent.

(4) Recall that the *binomial coefficient* $\binom{m}{k}$ is given by

$$\binom{m}{k} = \frac{m!}{k!(m-k)!},$$

with $0 \leq k \leq m$.

For any $m \geq 1$, we have the $m+1$ *Bernstein polynomials* of degree m given by

$$B_k^m(t) = \binom{m}{k} (1-t)^{m-k} t^k, \quad 0 \leq k \leq m.$$

Prove that

$$B_k^m(t) = \sum_{j=k}^m (-1)^{j-k} \binom{m}{j} \binom{j}{k} t^j. \quad (*)$$

Use the above to prove that $B_0^m(t), \dots, B_m^m(t)$ are linearly independent.

(5) Prove that

$$B_0^m(t) + \dots + B_m^m(t) = 1,$$

Extra credit (20 pts). What can you say about the symmetries of the $(m+1) \times (m+1)$ matrix expressing B_0^m, \dots, B_m^m in terms of the basis $1, t, \dots, t^m$?

Prove your claim (beware that in equation $(*)$ the coefficient of t^j in B_k^m is the entry on the $(k+1)$ th row of the $(j+1)$ th column, since $0 \leq k, j \leq m$. Make appropriate modifications to the indices).

What can you say about the sum of the entries on each row of the above matrix? What about the sum of the entries on each column?

(6) The purpose of this question is to express the t^i in terms of the Bernstein polynomials $B_0^m(t), \dots, B_m^m(t)$, with $0 \leq i \leq m$.

First, prove that

$$t^i = \sum_{j=0}^{m-i} t^i B_j^{m-i}(t), \quad 0 \leq i \leq m.$$

Then prove that

$$\binom{m}{i} \binom{m-i}{j} = \binom{m}{i+j} \binom{i+j}{i}.$$

Use the above facts to prove that

$$t^i = \sum_{j=0}^{m-i} \frac{\binom{i+j}{i}}{\binom{m}{i}} B_{i+j}^m(t).$$

Conclude that the Bernstein polynomials $B_0^m(t), \dots, B_m^m(t)$ form a basis of the vector space of polynomials of degree $\leq m$.

Compute the matrix expressing $1, t, t^2$ in terms of $B_0^2(t), B_1^2(t), B_2^2(t)$, and the matrix expressing $1, t, t^2, t^3$ in terms of $B_0^3(t), B_1^3(t), B_2^3(t), B_3^3(t)$.

You should find

$$\begin{pmatrix} 1 & 1 & 1 \\ 0 & 1/2 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

and

$$\begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1/3 & 2/3 & 1 \\ 0 & 0 & 1/3 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

(7) A *polynomial curve* $C(t)$ of degree m in the plane is the set of points $C(t) = \begin{pmatrix} x(t) \\ y(t) \end{pmatrix}$ given by two polynomials of degree $\leq m$,

$$\begin{aligned} x(t) &= \alpha_0 t^{m_1} + \alpha_1 t^{m_1-1} + \dots + \alpha_{m_1} \\ y(t) &= \beta_0 t^{m_2} + \beta_1 t^{m_2-1} + \dots + \beta_{m_2}, \end{aligned}$$

with $1 \leq m_1, m_2 \leq m$ and $\alpha_0, \beta_0 \neq 0$.

Prove that there exist $m+1$ points $b_0, \dots, b_m \in \mathbb{R}^2$ so that

$$C(t) = \begin{pmatrix} x(t) \\ y(t) \end{pmatrix} = B_0^m(t)b_0 + B_1^m(t)b_1 + \dots + B_m^m(t)b_m$$

for all $t \in \mathbb{R}$, with $C(0) = b_0$ and $C(1) = b_m$. Are the points b_1, \dots, b_{m-1} generally on the curve?

We say that the curve C is a *Bézier curve* and (b_0, \dots, b_m) is the list of *control points* of the curve (control points need not be distinct).

Remark: Because $B_0^m(t) + \dots + B_m^m(t) = 1$ and $B_i^m(t) \geq 0$ when $t \in [0, 1]$, the curve segment $C[0, 1]$ corresponding to $t \in [0, 1]$ belongs to the convex hull of the control points. This is an important property of Bézier curves which is used in geometric modeling to find the intersection of curve segments. Bézier curves play an important role in computer graphics and geometric modeling, but also in robotics because they can be used to model the trajectories of moving objects.

Problem B5 (40 pts). (a) Let A be an $n \times n$ matrix. If A is invertible, prove that for any $x \in \mathbb{R}^n$, if $Ax = 0$, then $x = 0$.

The converse is true: If for all $x \in \mathbb{R}^n$, $Ax = 0$ implies that $x = 0$, then A is invertible. We will prove this fact later, and you may use it without proof in part (b) of this problem.

(b) Let A be an $m \times n$ matrix and let B be an $n \times m$ matrix. Prove that $I_m - AB$ is invertible iff $I_n - BA$ is invertible.

TOTAL: 210 + 20 points.