

the entries  $\lambda_1, \dots, \lambda_n$  in  $D$  (the eigenvalues of  $A$ ) have absolute value  $+1$ . Thus, the entries in  $D$  are of the form  $\cos \theta + i \sin \theta = e^{i\theta}$ . Thus, we can assume that  $D$  is a diagonal matrix of the form

$$D = \begin{pmatrix} e^{i\theta_1} & & \cdots & \\ & e^{i\theta_2} & & \\ \vdots & \vdots & \ddots & \vdots \\ & & \cdots & e^{i\theta_p} \end{pmatrix}.$$

If we let  $E$  be the diagonal matrix

$$E = \begin{pmatrix} i\theta_1 & & \cdots & \\ & i\theta_2 & & \\ \vdots & \vdots & \ddots & \vdots \\ & & \cdots & i\theta_p \end{pmatrix}$$

it is obvious that  $E$  is skew Hermitian and that

$$e^E = D.$$

Then, letting  $B = UEU^*$ , we have

$$e^B = A,$$

and it is immediately verified that  $B$  is skew Hermitian, since  $E$  is.

If  $A$  is a unitary matrix with determinant  $+1$ , since the eigenvalues of  $A$  are  $e^{i\theta_1}, \dots, e^{i\theta_p}$  and the determinant of  $A$  is the product

$$e^{i\theta_1} \cdots e^{i\theta_p} = e^{i(\theta_1 + \cdots + \theta_p)}$$

of these eigenvalues, we must have

$$\theta_1 + \cdots + \theta_p = 0,$$

and so,  $E$  is skew Hermitian and has zero trace. As above, letting

$$B = UEU^*,$$

we have

$$e^B = A,$$

where  $B$  is skew Hermitian and has null trace.  $\square$

We now extend the result of Section 14.3 to Hermitian matrices.

## 14.5 Hermitian Matrices, Hermitian Positive Definite Matrices, and the Exponential Map

Recall that a Hermitian matrix is called *positive* (or *positive semidefinite*) if its eigenvalues are all positive or null, and *positive definite* if its eigenvalues

are all strictly positive. We denote the real vector space of Hermitian  $n \times n$  matrices by  $\mathbf{H}(n)$ , the set of Hermitian positive matrices by  $\mathbf{HP}(n)$ , and the set of Hermitian positive definite matrices by  $\mathbf{HPD}(n)$ .

The next lemma shows that every Hermitian positive definite matrix  $A$  is of the form  $e^B$  for some unique Hermitian matrix  $B$ . As in the real case, the set of Hermitian matrices is a real vector space, but it is not a Lie algebra because the Lie bracket  $[A, B]$  is not Hermitian unless  $A$  and  $B$  commute, and the set of Hermitian (positive) definite matrices is not a multiplicative group.

**Lemma 14.5.1** *For every Hermitian matrix  $B$ , the matrix  $e^B$  is Hermitian positive definite. For every Hermitian positive definite matrix  $A$ , there is a unique Hermitian matrix  $B$  such that  $A = e^B$ .*

*Proof.* It is basically the same as the proof of Theorem 14.5.1, except that a Hermitian matrix can be written as  $A = UDU^*$ , where  $D$  is a real diagonal matrix and  $U$  is unitary instead of orthogonal.  $\square$

Lemma 14.5.1 can be reformulated as stating that the map  $\exp: \mathbf{H}(n) \rightarrow \mathbf{HPD}(n)$  is a bijection. In fact, it can be shown that it is a homeomorphism. In the case of complex invertible matrices, the polar form theorem can be reformulated as stating that there is a bijection between the topological space  $\mathbf{GL}(n, \mathbb{C})$  of complex  $n \times n$  invertible matrices (also a group) and  $\mathbf{U}(n) \times \mathbf{HPD}(n)$ . As a corollary of the polar form theorem and Lemma 14.5.1, we have the following result: For every complex invertible matrix  $A$ , there is a unique unitary matrix  $U$  and a unique Hermitian matrix  $S$  such that

$$A = U e^S.$$

Thus, we have a bijection between  $\mathbf{GL}(n, \mathbb{C})$  and  $\mathbf{U}(n) \times \mathbf{H}(n)$ . But  $\mathbf{H}(n)$  itself is isomorphic to  $\mathbb{R}^{n^2}$ , and so there is a bijection between  $\mathbf{GL}(n, \mathbb{C})$  and  $\mathbf{U}(n) \times \mathbb{R}^{n^2}$ . It can also be shown that this bijection is a homeomorphism. This is an interesting fact. Indeed, this homeomorphism essentially reduces the study of the topology of  $\mathbf{GL}(n, \mathbb{C})$  to the study of the topology of  $\mathbf{U}(n)$ . This is nice, since it can be shown that  $\mathbf{U}(n)$  is compact (as a real manifold).

In the polar decomposition  $A = Ue^S$ , we have  $|\det(U)| = 1$ , since  $U$  is unitary, and  $\operatorname{tr}(S)$  is real, since  $S$  is Hermitian (since it is the sum of the eigenvalues of  $S$ , which are real), so that  $\det(e^S) > 0$ . Thus, if  $\det(A) = 1$ , we must have  $\det(e^S) = 1$ , which implies that  $S \in \mathbf{H}(n) \cap \mathfrak{sl}(n, \mathbb{C})$ . Thus, we have a bijection between  $\mathbf{SL}(n, \mathbb{C})$  and  $\mathbf{SU}(n) \times (\mathbf{H}(n) \cap \mathfrak{sl}(n, \mathbb{C}))$ .

In the next section we study the group  $\mathbf{SE}(n)$  of affine maps induced by orthogonal transformations, also called rigid motions, and its Lie algebra. We will show that the exponential map is surjective. The groups  $\mathbf{SE}(2)$  and  $\mathbf{SE}(3)$  play a fundamental role in robotics, dynamics, and motion planning.