

PRINT NAME: _____

LGIC 010 & PHIL 005
Practice Final Examination
Spring Term, 2009

1. Let S be the schema $(\forall x)\neg Lxx$.

- (a) (10 points) How long a list of distinct structures with universe of discourse $\{1, 2, 3\}$ satisfy the schema S ?
- (b) (10 points) How long a list of pairwise nonisomorphic structures with universe of discourse $\{1, 2, 3\}$ satisfy the schema S ?
- (c) (10 points) Give an example of a structure A with the following properties:
- $A \models S$;
 - $U^A = \{1, 2, 3\}$;
 - A has exactly three automorphisms;
 - exactly two subsets of $\{1, 2, 3\}$ are definable in A .

$L^A =$

2. (70 pts.) For each of the following pairs consisting of a set of schemata X and a schema S determine whether X implies S . If so, provide a deduction to establish the implication. If not, specify a structure which makes S false and all the schemata in X true.

(a) $X : \{(\exists x)(Px \vee Qx)\}$
 $S : (\exists x)Px \vee (\exists x)Qx$

$$A : U^A =$$

$$P^A =$$

$$Q^A =$$

Deduction

$$(b) \quad X : \{(\forall x)(Px \vee Qx)\}$$

$$S : (\forall x)Px \vee (\forall x)Qx$$

$$B : U^B =$$

$$P^B =$$

$$Q^B =$$

Deduction

$$(c) \quad X : \{(\forall x)Rxx, \neg(\forall x)(\forall y)Rxy\}$$

$$S : \neg(\exists x)(\forall y)x = y$$

$$C : U^C =$$

$$R^C =$$

Deduction

- (d) $X : \{(\forall x)(Fx \supset (\exists y)(\neg Fy \wedge (\forall z)(Rxz \equiv z = y))), (\forall x)(\neg Fx \supset (\exists y)(Fy \wedge (\forall z)(Rzx \equiv z = y))), (\forall x)(\forall y)(\forall z)((Pxy \wedge Pxz) \supset y = z), (\forall x)(\exists y)(Fy \wedge Pyx)\}$
 $S : p \wedge \neg p$

$$D : U^D =$$

$$F^D =$$

$$R^D =$$

$$P^D =$$

Deduction

(e) $X : \{(\forall x)(\forall y)(\forall z)((Rxy \wedge Ryz) \supset Rxz), (\forall x)\neg Rxx, (\forall x)(\forall y)(Rxy \vee Ryx \vee x = y), (\forall x)(\exists y)(\forall z)(Pxz \equiv z = y), (\forall x)(\exists y)Pyx, (\forall x)(\forall y)(Pxy \supset Rxy)\}$
 $S : p \wedge \neg p$

$E : U^E =$

$P^E =$

$R^E =$

Deduction

$$(f) \quad X : \{(\exists x)(\forall y)((\forall z)Rzy \equiv y = x)\}$$

$$S : (\forall x)(\exists y)(\forall z)(Rxz \equiv z = y)$$

$$F : U^F =$$

$$R^F =$$

Deduction

$$(g) \quad X : \{(\forall x)(\exists z)(\forall w)(Rwx \equiv w = z), \\ (\forall x)(\forall y)(\forall z)((Rxz \wedge Ryz) \supset y = x)\} \\ S : (\forall z)(\exists x)Rxz$$

$$G : U^G =$$

$$R^G =$$

Deduction