

PRINT NAME: Scott Weinstein

LGIC 010 & PHIL 005
Practice Final Examination
Spring Term, 2011

1. Let S be the schema $(\forall x)\neg Lxx$.

- (a) (10 points) How long a list of distinct structures with universe of discourse $\{1, 2, 3\}$ satisfy the schema S . 64
- (b) (10 points) How long a list of pairwise nonisomorphic structures with universe of discourse $\{1, 2, 3\}$ satisfy the schema S . 16
- (c) (10 points) Give an example of a structure A with the following properties:
- $A \models S$;
 - $U^A = \{1, 2, 3\}$;
 - A has exactly three automorphisms;
 - exactly two subsets of $\{1, 2, 3\}$ are definable in A .

$$L^A = \{\langle 1, 2 \rangle, \langle 2, 3 \rangle, \langle 3, 1 \rangle\}$$

2. (70 pts.) For each of the following pairs consisting of a set of schemata X and a schema S determine whether X implies S . If so, provide a deduction to establish the implication. If not, specify a structure which makes S false and all the schemata in X true.

$$(a) \quad X : \{(\exists x)(Px \vee Qx)\}$$

$$S : (\exists x)Px \vee (\exists x)Qx$$

$$A : U^A =$$

$$P^A =$$

$$Q^A =$$

Deduction

{1}	(1) $(\exists x)(Px \vee Qx)$	P
{1, 2}	(2) $Px \vee Qx$	(1) x EII
{3}	(3) Px	P
{4}	(4) Qx	P
{3}	(5) $(\exists x)Px$	(3) EG
{4}	(6) $(\exists x)Qx$	(4) EG
{}	(7) $Px \supset (\exists x)Px$	(5){3} D
{}	(8) $Qx \supset (\exists x)Qx$	(6){4} D
{1, 2}	(9) $(\exists x)Px \vee (\exists x)Qx$	(2)(7)(8) TF
{1}	(10) $(\exists x)Px \vee (\exists x)Qx$	(9){2} EIE

$$(b) \quad X : \{(\forall x)(Px \vee Qx)\}$$

$$S : (\forall x)Px \vee (\forall x)Qx$$

$$B : U^B = \{0, 1\}$$

$$P^B = \{0\}$$

$$Q^B = \{1\}$$

Deduction

(c) $X : \{(\forall x)Rxx, \neg(\forall x)(\forall y)Rxy\}$
 $S : \neg(\exists x)(\forall y)x = y$

$C : U^C =$

$R^C =$

Deduction

{1}	(1) $(\forall x)Rxx$	P
{2}	(2) $\neg(\forall x)(\forall y)Rxy$	P
{3}	(3) $(\exists x)(\forall y)x = y$	P
{3, 4}	(4) $(\forall y)u = y$	(3)u EII
{1}	(5) Ruu	(1) UI
{3, 4}	(6) $u = y$	(4) UI
{}	(7) $u = y \supset (Ruu \equiv Ruy)$	III
{3, 4}	(8) $u = x$	(4) UI
{}	(9) $u = x \supset (Ruy \equiv Rxy)$	III
{1, 3, A}	(10) Rxy	(5)(6) TF; (7)(8) {4} EIE (9)
{1, 3}	(11) $(\forall y)Rxy$	(10) UG
{1, 3}	(12) $(\forall x)(\forall y)Rxy$	(11) UG
{1, 2, 3}	(13) $p \wedge \neg p$	(2)(12) TF
{1, 2}	(14) $(\exists x)(\forall y)x = y \supset (p \wedge \neg p)$	{3}(13) D
{1, 2}	(15) $\neg(\exists x)(\forall y)x = y$	(14) TF

(d) $X : \{(\forall x)(Fx \supset (\exists y)(\neg Fy \wedge (\forall z)(Rxz \equiv z = y))), (\forall x)(\neg Fx \supset (\exists y)(Fy \wedge (\forall z)(Rzx \equiv z = y))), (\forall x)(\forall y)(\forall z)((Pxy \wedge Pxz) \supset y = z), (\forall x)(\exists y)(Fy \wedge Pyx)\}$
 $S : p \wedge \neg p$

$$D : U^D = \{0, 1, 2, \dots\}$$

$$F^D = \{0, 2, 4, \dots\}$$

$$R^D = \{\langle 2i, 2i + 1 \rangle \mid 0 \leq i\}$$

$$P^D = \{\langle 2i, i \rangle \mid 0 \leq i\}$$

Deduction

(e) $X : \{(\forall x)(\forall y)(\forall z)((Rxy \wedge Ryz) \supset Rxz), (\forall x)\neg Rxx, (\forall x)(\forall y)(Rxy \vee Ryx \vee x = y), (\forall x)(\exists y)(\forall z)(Pxz \equiv z = y), (\forall x)(\exists y)Pyx, (\forall x)(\forall y)(Pxy \supset Rxy)\}$
 $S : p \wedge \neg p$

$$E : U^E = \{\dots -3, -2, -1, 0, 1, 2, 3, \dots\}$$

$$P^E = \{\langle i, i+1 \rangle \mid i \in U^E\}$$

$$R^E = \{\langle i, j \rangle \mid i < j\}$$

Deduction

$$(f) \quad X : \{(\exists x)(\forall y)((\forall z)Rzy \equiv y = x)\}$$

$$S : (\forall x)(\exists y)(\forall z)(Rxz \equiv z = y)$$

$$F : U^F = \{0, 1\}$$

$$R^F = \{\langle 0, 0 \rangle, \langle 0, 1 \rangle, \langle 1, 0 \rangle\}$$

Deduction

$$(g) \quad X : \{(\forall x)(\exists z)(\forall w)(Rwx \equiv w = z), \\ (\forall x)(\forall y)(\forall z)((Rxz \wedge Ryz) \supset y = x)\} \\ S : (\forall z)(\exists x)Rxz$$

$$G : U^G = \{0, 1, 2, 3, \dots\}$$

$$R^G = \{\langle i, i + 1 \rangle \mid i \in U^G\}$$

Deduction