

PRINT NAME: \_\_\_\_\_

**LGIC 010 & PHIL 005**  
**Practice Final Examination**  
**Spring Term, 2012**

1. Let  $S$  be the schema

$$(\forall x)(\forall y)(Lxy \supset Lyx) \wedge (\forall x)\neg Lxx.$$

- (a) (10 points) How long a list of distinct structures with universe of discourse  $\{1, 2, 3, 4\}$  satisfy the schema  $S$ ?
- (b) (10 points) How long a list of pairwise nonisomorphic structures with universe of discourse  $\{1, 2, 3, 4\}$  satisfy the schema  $S$ ?
- (c) (10 points) Give an example of a structure  $A$  with the following properties:
- $A \models S$ ;
  - $U^A = \{1, 2, 3, 4\}$ ;
  - $A$  has exactly two automorphisms;
  - exactly eight subsets of  $\{1, 2, 3, 4\}$  are definable in  $A$ .

$$L^A =$$

2. (70 pts.) For each of the following pairs consisting of a set of schemata  $X$  and a schema  $S$  determine whether  $X$  implies  $S$ . If so, provide a deduction to establish the implication. If not, specify a structure which makes  $S$  false and all the schemata in  $X$  true.

(a)  $X : \{(\exists x)Px \wedge (\exists x)Qx\}$   
 $S : (\exists x)(Px \wedge Qx)$

$$A : U^A =$$

$$P^A =$$

$$Q^A =$$

Deduction

$$(b) \quad X : \{(\forall x)(Px \wedge Qx)\}$$

$$S : (\forall x)Px \wedge (\forall x)Qx$$

$$B : U^B =$$

$$P^B =$$

$$Q^B =$$

Deduction

(c)  $X : \{(\forall x)((\exists y)Lxy \supset (\forall z)Lzx), (\exists x)(\exists y)Lxy\}$   
 $S : (\forall u)(\forall z)Lzu$

$C : U^C =$

$L^C =$

Deduction

$$(d) \quad X : \{(\exists x)(\forall y)Rxy\}$$

$$S : (\forall x)(\exists y)Rxy$$

$$D : U^D =$$

$$R^D =$$

Deduction

(e)  $X : \{(\forall x)\neg Lxx, (\forall x)(\forall y)(Lxy \supset Lyx),$   
 $(\forall x)(\exists y)(\exists z)(Lyz \wedge (\forall w)(Lxw \equiv (w = y \vee w = z))),$   
 $(\forall v)(\forall w)(\forall x)(\forall y)(\forall z)((Rvwz \wedge Rxyz) \supset (v = x \wedge w = y)),$   
 $(\forall x)(\forall y)(\forall z)(Rxyz \supset (Px \wedge Py)), (\forall z)(\exists x)(\exists y)Rxyz,$   
 $(\forall x)(\forall y)((Px \wedge Py) \supset (\exists z)(\forall w)(Rxyw \equiv w = z))\}$   
 $S : (\exists x)(\exists y)(Px \wedge Py \wedge Lxy)$

$E : U^E =$

$L^E =$

$P^E =$

$R^E =$

Deduction

$$(f) \quad X : \{(\exists x)(\forall y)(Fy \equiv x = y)\}$$

$$S : (\exists x)Fx \wedge (\forall x)(\forall y)((Fx \wedge Fy) \supset x = y)$$

$$F : U^F =$$

$$F^F =$$

Deduction

$$(g) \quad X : \{(\forall x)(\exists z)(\forall w)(Rwx \equiv w = z), (\forall z)(\exists x)Rxz\}$$

$$S : (\forall x)(\forall y)(\forall z)((Rxz \wedge Ryz) \supset y = x)$$

$$G : U^G =$$

$$R^G =$$

Deduction