

**LGIC 010 & PHIL 005**  
**Problem Set 8**  
**Spring Term, 2016**  
**DUE IN CLASS MONDAY, APRIL 11**

- $\mathbb{Z}$  is the set of integers  $\{\dots - 3, -2, -1, 0, 1, 2, 3, \dots\}$ .
- $\mathbb{Z}^+$  is the set of positive integers  $\{1, 2, 3, \dots\}$ .
- A positive integer  $i$  is a *prime number* if and only if  $i > 1$  and the only divisors of  $i$  are  $i$  and 1. The first few prime numbers are 2, 3, 5, 7, 11,  $\dots$
- Let  $A$  be a structure interpreting a triadic predicate letter  $P$ .  $h$  is an *automorphism* of  $A$  if and only if  $h$  is a bijection of  $U^A$  onto  $U^A$  and for all  $i, j, k \in U^A$ ,  $\langle i, j, k \rangle \in P^A$  if and only if  $\langle h(i), h(j), h(k) \rangle \in P^A$ .
- Let  $A$  be a structure.  $\text{Aut}(A) = \{h \mid h \text{ is an automorphism of } A\}$ .
- Let  $S(x)$  be a schema with a single free variable  $x$  and  $A$  a structure.

$$S[A] = \{a \in U^A \mid A \models S[x|a]\}.$$

1. Let  $A_1$  be a structure interpreting a single triadic predicate letter  $P_1$  with  $U^A = \mathbb{Z}^+$  and  $P_1^A = \{\langle i, j, k \rangle \mid i \cdot j = k\}$ , the product relation on  $\mathbb{Z}^+$ .
  - (a) Let  $X_1 = \{i \in \mathbb{Z}^+ \mid i \text{ is a prime number}\}$ . Is  $X_1$  definable in  $A_1$ ? If it is, write down a schema  $S_1(x)$  such that  $S_1[A_1] = X_1$ ; if not, specify an  $h_1 \in \text{Aut}(A_1)$  such that  $h_1[X_1] \neq X_1$ .
  - (b) Let  $X_2 = \{i \in \mathbb{Z}^+ \mid i \text{ is a positive power of } 2\} = \{2, 4, 8, 16, \dots\}$ . Is  $X_2$  definable in  $A_1$ ? If it is, write down a schema  $S_2(x)$  such that  $S_2[A_1] = X_2$ ; if not, specify an  $h_2 \in \text{Aut}(A_1)$  such that  $h_2[X_2] \neq X_2$ .
  - (c) Let  $X_3 = \{i \in \mathbb{Z}^+ \mid i \text{ is a positive power of a prime number}\} = \{2, 4, \dots, 3, 9, \dots\}$ . Is  $X_3$  definable in  $A_1$ ? If it is, write down a schema  $S_3(x)$  such that  $S_3[A_1] = X_3$ ; if not, specify an  $h_3 \in \text{Aut}(A_1)$  such that  $h_3[X_3] \neq X_3$ .
2. Let  $A_2$  be a structure interpreting a single triadic predicate letter  $P_2$  with  $U^A = \mathbb{Z}$  and  $P_2^A = \{\langle i, j, k \rangle \mid i + j = k\}$ , the sum relation on  $\mathbb{Z}$ .
  - (a) Let  $X_4 = \{i \in \mathbb{Z} \mid i \neq 0\}$ . Is  $X_4$  definable in  $A_2$ ? If it is, write down a schema  $S_4(x)$  such that  $S_4[A_2] = X_4$ ; if not, specify an  $h_4 \in \text{Aut}(A_2)$  such that  $h_4[X_4] \neq X_4$ .
  - (b) Let  $X_5 = \{i \in \mathbb{Z} \mid i > 0\}$ . Is  $X_5$  definable in  $A_2$ ? If it is, write down a schema  $S_5(x)$  such that  $S_5[A_2] = X_5$ ; if not, specify an  $h_5 \in \text{Aut}(A_2)$  such that  $h_5[X_5] \neq X_5$ .
  - (c) Let  $X_6 = \{i \in \mathbb{Z} \mid i \text{ is divisible by } 3 \text{ (without remainder)}\}$ . Is  $X_6$  definable in  $A_2$ ? If it is, write down a schema  $S_6(x)$  such that  $S_6[A_2] = X_6$ ; if not, specify an  $h_6 \in \text{Aut}(A_2)$  such that  $h_6[X_6] \neq X_6$ .