

Convergence of No-Regret Play to Nash Equilibrium

Aaron Roth

University of Pennsylvania

February 15 2024

Overview

- ▶ We've seen that two-player zero sum games are special.

Overview

- ▶ We've seen that two-player zero sum games are special.
- ▶ They have a *value*, order of play doesn't matter, equilibria can be computed "easily"

Overview

- ▶ We've seen that two-player zero sum games are special.
- ▶ They have a *value*, order of play doesn't matter, equilibria can be computed "easily"
- ▶ i.e. it does not require counterspeculation — don't need to reason about your opponent to compute a minmax strategy.

Overview

- ▶ We've seen that two-player zero sum games are special.
- ▶ They have a *value*, order of play doesn't matter, equilibria can be computed "easily"
- ▶ i.e. it does not require counterspeculation — don't need to reason about your opponent to compute a minmax strategy.
- ▶ But you need to understand the game extremely well and make careful calculations.

Overview

- ▶ We've seen that two-player zero sum games are special.
- ▶ They have a *value*, order of play doesn't matter, equilibria can be computed "easily"
- ▶ i.e. it does not require counterspeculation — don't need to reason about your opponent to compute a minmax strategy.
- ▶ But you need to understand the game extremely well and make careful calculations.
- ▶ Is there a natural dynamic that leads to Nash equilibrium if everyone uses it?

Overview

- ▶ We've seen that two-player zero sum games are special.
- ▶ They have a *value*, order of play doesn't matter, equilibria can be computed "easily"
- ▶ i.e. it does not require counterspeculation — don't need to reason about your opponent to compute a minmax strategy.
- ▶ But you need to understand the game extremely well and make careful calculations.
- ▶ Is there a natural dynamic that leads to Nash equilibrium if everyone uses it?
- ▶ How many of these properties depend on the "two player" caveat?

Two players?

Do these special properties carry over to general n player zero sum games?

Two players?

Do these special properties carry over to general n player zero sum games? We can certainly define such games:

Definition

An n player game is zero-sum if for every action profile $a \in A$,
$$\sum_{i=1}^n u_i(a) = 0.$$

Two players?

Do these special properties carry over to general n player zero sum games? We can certainly define such games:

Definition

An n player game is zero-sum if for every action profile $a \in A$,
$$\sum_{i=1}^n u_i(a) = 0.$$

The answer is no.

“Meta Theorem”: n player zero-sum games don't have any special properties that $n - 1$ player general sum games don't have.

In particular, we should not expect such games to have a value, nor that their equilibria should be easy to compute.

Two players?

Do these special properties carry over to general n player zero sum games? We can certainly define such games:

Definition

An n player game is zero-sum if for every action profile $a \in A$, $\sum_{i=1}^n u_i(a) = 0$.

The answer is no.

“Meta Theorem”: n player zero-sum games don't have any special properties that $n - 1$ player general sum games don't have.

In particular, we should not expect such games to have a value, nor that their equilibria should be easy to compute.

“Proof”: Any $n - 1$ player game can be made into an n player zero sum game, by adding a new player n (with a trivial action set), and $u_n(a) = -\sum_{i=1}^{n-1} u_i(a)$. Since player n is payoff irrelevant to the $n - 1$ other players, the equilibrium structure remains identical to the original game.

But we can generalize with more structure...

Definition

A separable graphical game is defined by a graph $G = (V, E)$. The set of players corresponds to the set of vertices: $P = V$. Each player's utility function is decomposable as a sum of neighbor-specific utility functions, one for each of his neighbors in G :

$$u_i(a) = \sum_{(i,j) \in E} u_i^{(i,j)}(a_i, a_j)$$

i.e. it is as if each player is playing a 2-player game with each of his neighbors – except he must pick a single action a_i to play simultaneously against each of his neighbors.

Separable Graphical Games

Separable Graphical Games

Zero sum separable graphical games have many of the properties of two player zero sum games:

Separable Graphical Games

Zero sum separable graphical games have many of the properties of two player zero sum games:

1. They continue to have a *value*

Separable Graphical Games

Zero sum separable graphical games have many of the properties of two player zero sum games:

1. They continue to have a *value*
2. Equilibria are easy to compute with efficient dynamics.

Separable Graphical Games

Zero sum separable graphical games have many of the properties of two player zero sum games:

1. They continue to have a *value*
2. Equilibria are easy to compute with efficient dynamics.
3. We don't require each of the constituent 2-player games are zero sum — just that the aggregate is.

Regret

Definition

A sequence of action profiles a^1, \dots, a^T has regret $\Delta(T)$ if for all players i and actions a_i^* we have:

$$\frac{1}{T} \sum_{t=1}^T u_i(a^t) \geq \frac{1}{T} \sum_{t=1}^T u_i(a_i^*, a_{-i}^t) - \Delta(T)$$

We say that such an action sequence is *no-regret* if $\Delta(T) = o_T(1)$.

Regret

Definition

A sequence of action profiles a^1, \dots, a^T has regret $\Delta(T)$ if for all players i and actions a_i^* we have:

$$\frac{1}{T} \sum_{t=1}^T u_i(a^t) \geq \frac{1}{T} \sum_{t=1}^T u_i(a_i^*, a_{-i}^t) - \Delta(T)$$

We say that such an action sequence is *no-regret* if $\Delta(T) = o_T(1)$.

1. How to generate a sequence of no-regret play?

Regret

Definition

A sequence of action profiles a^1, \dots, a^T has regret $\Delta(T)$ if for all players i and actions a_i^* we have:

$$\frac{1}{T} \sum_{t=1}^T u_i(a^t) \geq \frac{1}{T} \sum_{t=1}^T u_i(a_i^*, a_{-i}^t) - \Delta(T)$$

We say that such an action sequence is *no-regret* if $\Delta(T) = o_T(1)$.

1. How to generate a sequence of no-regret play?
2. Have every player play polynomial weights. Then

$$\Delta(T) = O\left(2\sqrt{\frac{\log k}{T}}\right)$$

Regret

Definition

A sequence of action profiles a^1, \dots, a^T has regret $\Delta(T)$ if for all players i and actions a_i^* we have:

$$\frac{1}{T} \sum_{t=1}^T u_i(a^t) \geq \frac{1}{T} \sum_{t=1}^T u_i(a_i^*, a_{-i}^t) - \Delta(T)$$

We say that such an action sequence is *no-regret* if $\Delta(T) = o_T(1)$.

1. How to generate a sequence of no-regret play?
2. Have every player play polynomial weights. Then

$$\Delta(T) = O\left(2\sqrt{\frac{\log k}{T}}\right)$$

3. But not the only way...

Regret

Definition

A sequence of action profiles a^1, \dots, a^T has regret $\Delta(T)$ if for all players i and actions a_i^* we have:

$$\frac{1}{T} \sum_{t=1}^T u_i(a^t) \geq \frac{1}{T} \sum_{t=1}^T u_i(a_i^*, a_{-i}^t) - \Delta(T)$$

We say that such an action sequence is *no-regret* if $\Delta(T) = o_T(1)$.

1. How to generate a sequence of no-regret play?
2. Have every player play polynomial weights. Then

$$\Delta(T) = O\left(2\sqrt{\frac{\log k}{T}}\right)$$

3. But not the only way...
4. A permissive *family* of dynamics.

Dynamics

Given a sequence of action profiles a^1, \dots, a^T , write $\bar{a}_i = \frac{1}{T} \sum_{t=1}^T a_i^t$ to denote the mixed strategy for player i that selects an action in $\{a_i^1, \dots, a_i^T\}$ uniformly at random.

Dynamics

Given a sequence of action profiles a^1, \dots, a^T , write $\bar{a}_i = \frac{1}{T} \sum_{t=1}^T a_i^t$ to denote the mixed strategy for player i that selects an action in $\{a_i^1, \dots, a_i^T\}$ uniformly at random.

Theorem

Consider any zero sum separable graphical game G . If a sequence of action profiles a^1, \dots, a^T has regret $\Delta(T)$, then the mixed strategies:

$$(\bar{a}_1, \dots, \bar{a}_n)$$

forms an $n\Delta(T)$ -approximate Nash equilibrium.

Dynamics

Given a sequence of action profiles a^1, \dots, a^T , write $\bar{a}_i = \frac{1}{T} \sum_{t=1}^T a_i^t$ to denote the mixed strategy for player i that selects an action in $\{a_i^1, \dots, a_i^T\}$ uniformly at random.

Theorem

Consider any zero sum separable graphical game G . If a sequence of action profiles a^1, \dots, a^T has regret $\Delta(T)$, then the mixed strategies:

$$(\bar{a}_1, \dots, \bar{a}_n)$$

forms an $n\Delta(T)$ -approximate Nash equilibrium.

If every player plays using polynomial weights, they converge to an ϵ -approximate Nash equilibrium by in:

$$T = \frac{4n^2 \log k}{\epsilon^2}$$

many rounds. In a two player game this is $T = 16 \log(k)/\epsilon^2$ steps.

Proof

1. A useful fact: for every action $a_i^* \in A_i$ we have:

$$\begin{aligned} \frac{1}{T} \sum_{t=1}^T \sum_{(i,j) \in E} u_i^{i,j}(a_i^*, a_j^t) &= \sum_{(i,j) \in E} \sum_{t=1}^T \frac{1}{T} u_i^{i,j}(a_i^*, a_j^t) \\ &= \sum_{(i,j) \in E} u_i^{i,j}(a_i^*, \bar{a}_j) \end{aligned}$$

Proof

1. A useful fact: for every action $a_i^* \in A_i$ we have:

$$\begin{aligned} \frac{1}{T} \sum_{t=1}^T \sum_{(i,j) \in E} u_i^{i,j}(a_i^*, a_j^t) &= \sum_{(i,j) \in E} \sum_{t=1}^T \frac{1}{T} u_i^{i,j}(a_i^*, a_j^t) \\ &= \sum_{(i,j) \in E} u_i^{i,j}(a_i^*, \bar{a}_j) \end{aligned}$$

2. Suppose every player i is playing according to \bar{a}_i . Let a_i^* be the best response of player i to the distribution of his opponents. We know:

$$\sum_{(i,j) \in E} u_i^{i,j}(a_i^*, \bar{a}_j) \geq \sum_{(i,j) \in E} u_i^{i,j}(\bar{a}_i, \bar{a}_j)$$

Proof

1. We also know, since a^1, \dots, a^t have $\Delta(T)$ regret, that for all $i \in P$:

$$\underbrace{\frac{1}{T} \sum_{t=1}^T \sum_{(i,j) \in E} u_i^{(i,j)}(a_i^t, a_j^t)}_{LHS} \geq \underbrace{\sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) - \Delta(T)}_{RHS}$$

Proof

1. We also know, since a^1, \dots, a^t have $\Delta(T)$ regret, that for all $i \in P$:

$$\underbrace{\frac{1}{T} \sum_{t=1}^T \sum_{(i,j) \in E} u_i^{(i,j)}(a_i^t, a_j^t)}_{LHS} \geq \underbrace{\sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) - \Delta(T)}_{RHS}$$

2. Summing the LHS over all players:

$$LHS = \frac{1}{T} \sum_{t=1}^T \sum_{i=1}^n \sum_{(i,j) \in E} u_i^{(i,j)}(a_i^t, a_j^t) = \frac{1}{T} \sum_{t=1}^T 0 = 0$$

(why?)

Proof

1. For all $i \in P$:

$$\underbrace{\frac{1}{T} \sum_{t=1}^T \sum_{(i,j) \in E} u_i^{(i,j)}(a_i^t, a_j^t)}_{LHS} \geq \underbrace{\sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) - \Delta(T)}_{RHS}$$

Proof

1. For all $i \in P$:

$$\underbrace{\frac{1}{T} \sum_{t=1}^T \sum_{(i,j) \in E} u_i^{(i,j)}(a_i^t, a_j^t)}_{LHS} \geq \underbrace{\sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) - \Delta(T)}_{RHS}$$

2. Now summing the RHS:

$$RHS = \sum_{i=1}^n \sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) - n \cdot \Delta(T)$$

Proof

1. Combining the bounds (LHS $>$ RHS):

Proof

1. Combining the bounds (LHS > RHS):
- 2.

$$n\Delta(T) \geq \sum_{i=1}^n \sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j)$$

Proof

1. Combining the bounds (LHS > RHS):
- 2.

$$\begin{aligned} n\Delta(T) &\geq \sum_{i=1}^n \sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) \\ &= \sum_{i=1}^n \left(\sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) - \sum_{(i,j) \in E} u_i^{i,j}(\bar{a}_i, \bar{a}_j) \right) \end{aligned}$$

Proof

1. Combining the bounds (LHS > RHS):
- 2.

$$\begin{aligned} n\Delta(T) &\geq \sum_{i=1}^n \sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) \\ &= \sum_{i=1}^n \left(\sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) - \sum_{(i,j) \in E} u_i^{i,j}(\bar{a}_i, \bar{a}_j) \right) \end{aligned}$$

3. (why?)

Proof

1. Combining the bounds (LHS > RHS):
- 2.

$$\begin{aligned} n\Delta(T) &\geq \sum_{i=1}^n \sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) \\ &= \sum_{i=1}^n \left(\sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) - \sum_{(i,j) \in E} u_i^{i,j}(\bar{a}_i, \bar{a}_j) \right) \end{aligned}$$

3. (why?)
4. Lets think about each term...

Proof

$$n\Delta(T) \geq \sum_{i=1}^n \left(\sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) - \sum_{(i,j) \in E} u_i^{i,j}(\bar{a}_i, \bar{a}_j) \right)$$

Proof

$$n\Delta(T) \geq \sum_{i=1}^n \left(\sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) - \sum_{(i,j) \in E} u_i^{i,j}(\bar{a}_i, \bar{a}_j) \right)$$

1. For each term we have:

$$\left(\sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) - \sum_{(i,j) \in E} u_i^{i,j}(\bar{a}_i, \bar{a}_j) \right) \geq 0$$

(why?)

Proof

$$n\Delta(T) \geq \sum_{i=1}^n \left(\sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) - \sum_{(i,j) \in E} u_i^{i,j}(\bar{a}_i, \bar{a}_j) \right)$$

1. For each term we have:

$$\left(\sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) - \sum_{(i,j) \in E} u_i^{i,j}(\bar{a}_i, \bar{a}_j) \right) \geq 0$$

(why?)

2. So for each player i :

$$\sum_{(i,j) \in E} u_i^{i,j}(\bar{a}_i, \bar{a}_j) \geq \sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) - n\Delta(T)$$

(why?)

Proof

$$n\Delta(T) \geq \sum_{i=1}^n \left(\sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) - \sum_{(i,j) \in E} u_i^{i,j}(\bar{a}_i, \bar{a}_j) \right)$$

1. For each term we have:

$$\left(\sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) - \sum_{(i,j) \in E} u_i^{i,j}(\bar{a}_i, \bar{a}_j) \right) \geq 0$$

(why?)

2. So for each player i :

$$\sum_{(i,j) \in E} u_i^{i,j}(\bar{a}_i, \bar{a}_j) \geq \sum_{(i,j) \in E} u_i^{(i,j)}(a_i^*, \bar{a}_j) - n\Delta(T)$$

(why?)

3. Tada!

Thanks!

See you next class — stay healthy!