

# Mathematical Foundations of Computer Science

## Lecture Outline

November 17, 2020

---

### Matching in Bipartite Graphs

An *independent set* of a graph is a set of pair-wise non-adjacent vertices. A *bipartite graph*,  $(U, V, E)$ , is a graph whose vertex set is  $U \cup V$  and for each edge  $e = (u, v) \in E$ ,  $u \in U$  and  $v \in V$ . In other words,  $U$  and  $V$  are independent sets and each edge in  $E$  connects a vertex in  $U$  to a vertex in  $V$ .

Now consider the following scenario. There is a set of girls and a set of boys. Each girl likes some boys and dislikes others. What conditions would guarantee that each girl is paired-up with a boy that she likes and that no two girls are paired-up with the same boy.

We can model this situation using a bipartite graph,  $(X, Y, E)$ , where each vertex in  $X$  represents a girl, each vertex in  $Y$  represents a boy and an edge  $(g, b) \in E$  means that girl  $g$  likes boy  $b$ . We are interested in the conditions that would guarantee a matching that saturates every vertex in  $X$ .

Hall's theorem gives the necessary and sufficient conditions for the existence of such matchings in bipartite graphs.

---

**Example. [Hall's Theorem]** Let  $G = (X, Y, E)$  be a bipartite graph. For any set  $S$  of vertices, let  $N_G(S)$  be the set of vertices adjacent to vertices in  $S$ . Prove that  $G$  contains a matching that saturates every vertex in  $X$  iff  $|N_G(S)| \geq |S|, \forall S \subseteq X$ . The condition "For all  $S \subseteq X, |N(S)| \geq |S|$ " is called Hall's condition.

**Solution.** We prove that Hall's condition is necessary as follows. Suppose  $G$  contains a matching  $M$  that saturates every vertex in  $X$ . Let  $S$  be a subset of  $X$ . Since each vertex in  $S$  is matched under  $M$  to a distinct vertex in  $N_G(S)$ ,  $|N_G(S)| \geq |S|$ .

We will now prove the sufficiency of Hall's condition, i.e., if  $|N_G(S)| \geq |S|, \forall S \subseteq X$  then  $G$  contains a matching that saturates every vertex in  $X$ . We prove this by induction on the size of  $X$ .

Base Case:  $|X| = 1$ . If the only vertex in  $X$  is connected to at least one vertex in  $Y$  then clearly a matching exists.

Induction Hypothesis: Assume that Hall's condition is sufficient when  $|X| = j$ , for all  $j$  such that  $1 \leq j \leq k$ .

Induction Step: We want to prove that the sufficiency of Hall's condition when  $|X| = k + 1$ . Let  $G = (X, Y, E)$  be a graph with  $k + 1$  vertices in  $X$  such that  $\forall S \subseteq X, |N_G(S)| \geq |S|$ .

We consider the following two cases.

**Case I:** For every non-empty proper subset  $W \subset X$ ,  $|N_G(W)| > |W|$ . In this case, we pair-up an arbitrary vertex  $x \in X$  with one of its neighbors, say  $y \in Y$ . Now consider the subgraph  $G' = (X', Y', E')$ , where  $X' = X \setminus \{x\}$ ,  $Y' = Y \setminus \{y\}$ , and  $E' = E \setminus \{(x, y)\}$ . After the removal of  $y$ , the neighborhood of any subset,  $S' \subseteq X'$  in  $G'$  is at most one less than its neighborhood in  $G$ . But since  $|N_G(S')| > |S'|$ , after removal of  $y$ , it must be that  $|N_{G'}(S')| \geq |S'|$ . Thus, Hall's condition holds for  $G'$ . By induction hypothesis,  $G'$  contains a matching  $M'$  that saturates every vertex in  $X'$ . Hence,  $M' \cup \{(x, y)\}$  is a matching that saturates every vertex in  $X$ .

**Case II:** For some non-empty proper subset  $W \subset X$ ,  $|N(W)| = |W|$ . For all  $S' \subseteq W$ , we have  $N_G(S') \subseteq N_G(W)$ . Hence, Hall's condition holds for the subgraph induced by  $W \cup N(W)$ . By induction hypothesis, there is a matching  $M_1$  that matches every vertex in  $W$  to a vertex in  $N_G(W)$ . Note that  $M_1$  is a perfect matching. Consider the subgraph  $G' = (X', Y', E')$ , where  $X' = X \setminus W$ ,  $Y' = Y \setminus N(W)$ , and  $E'$  consists of all edges between  $X'$  and  $Y'$ . If we can prove that Hall's condition holds for  $G'$  then by induction hypothesis,  $G'$  has a matching  $M_2$  that saturates every vertex in  $X'$ . Then,  $M_1 \cup M_2$  is clearly a matching in  $G$  that saturates every vertex in  $X$ . It now remains to prove that  $\forall T \subseteq X', |N_{G'}(T)| \geq |T|$ . Note that  $N_G(W \cup T) = N_G(W) \cup N_{G'}(T)$ ,  $|N_G(W)| = |W|$ ,  $W$  and  $T$  are disjoint, and  $N_G(W)$  and  $N_{G'}(T)$  are disjoint. Then,

$$\begin{aligned} |N_G(W \cup T)| &\geq |W \cup T| \text{ (follows because } \forall S \subseteq X, |N_G(S)| \geq |S|) \\ |N_G(W)| + |N_{G'}(T)| &\geq |W| + |T| \\ |W| + |N_{G'}(T)| &\geq |W| + |T| \\ |N_{G'}(T)| &\geq |T| \end{aligned}$$

This proves the sufficiency of Hall's condition.

## Relations

A *binary relation* is a set of ordered pairs. For example, let  $R = \{(1, 2), (2, 3), (5, 4)\}$ . Then since  $(1, 2) \in R$ , we say that 1 is related to 2 by relation  $R$ . We denote this by  $1 R 2$ . Similarly, since  $(4, 7) \notin R$ , 4 is not related to 7 by relation  $R$ , denoted by  $4 \not R 7$ .

A binary relation  $R$  from set  $A$  to set  $B$  is a subset of the cartesian product  $A \times B$ . When  $A = B$ , we say that  $R$  is a relation on set  $A$ .

**Example.** Let  $A = \{1, 2, 3, 4\}$  and  $B = \{a, b, c\}$ . Consider the following relations.

$$\begin{aligned} R_1 &= \{(1, 1), (1, 2), (2, 2), (2, 3)\} \\ R_2 &= \{(1, 2), (2, 3), (3, 4), (4, 1), (4, 4)\} \\ R_3 &= \{(1, a), (2, a), (3, b), (4, c)\} \\ R_4 &= \{(a, 1), (a, 3), (a, 4), (c, 1)\} \\ R_5 &= \{(a, a), (a, b), (1, c)\} \end{aligned}$$

$R_1$  and  $R_2$  are relations on  $A$ .  $R_3$  is a relation from  $A$  to  $B$ .  $R_4$  is a relation from  $B$  to  $A$ .  $R_5$  is not a relation on sets  $A$  and  $B$  and it is neither a relation from  $A$  to  $B$  nor a relation from  $B$  to  $A$ .

Below are some more examples of relations.

- If  $S$  is a set then “is a subset of”,  $\subseteq$  is a relation on  $\mathcal{P}(S)$ , the power set of  $S$ .
- “is a student in” is a relation from the set of students to the set of courses.
- “=” is a relation on  $\mathbb{Z}$ .
- “has a path in  $G$  to” is a relation on  $V(G)$ , the set of vertices in  $G$ .

**Example.** How many relations are there on a set  $A$  of  $n$  elements?

**Solution.** Note that any relation on  $A$  is a subset of  $A \times A$  and since the power set of  $A \times A$  contains all subsets of  $A \times A$ , the number of possible relations on  $A$  is the cardinality of the power set of  $A \times A$ . Since  $|A \times A| = n^2$ , the cardinality of the power set of  $A \times A$  is  $2^{n^2}$ . Thus our answer is  $2^{n^2}$ .