

Contagion in Networks

Networked Life

NETS 112

Fall 2016

Prof. Michael Kearns


Two Models of Network Formation

- Start with a grid, remove random fraction of vertices
 - “local” or “geographic” connectivity
- Start with N isolated vertices, add random edges
 - “long distance” connectivity
- Examine a deterministic contagion model
- Widespread infection occurs at “tipping point” of connectivity

“Mathematizing” the Forest Fire

(see Coursera “Contagion” video)

- Start with a regular 2-dimensional grid network
 - this represents a complete forest
- Delete each vertex (and all 4 of its edges) with probability $1-p$
 - p is fraction of forest, $1-p$ is fraction of parking lots or clear-cut
- Choose a random remaining vertex v
 - this is my campsite
- Q: What is the expected size of v 's *connected component*?
 - i.e. the number of vertices reachable from v
 - this is how much of the forest is going to burn
- Observe a “tipping point” around $p = 0.6$



“Mathematizing” the Average Degree Demo (see Coursera “Contagion” video)

- Let d be the desired average degree in a network of N vertices
- Then the total number of edges should be

$$e = d \times N / 2$$


- Just start connecting random pairs of vertices until you have e edges
- Pick a random vertex v to infect
- What is the size of v 's connected component?
- Observe a “tipping point” around $d=3$

Some Remarks on the Demos


- Connectivity patterns were either *local* or *random*
 - will eventually formalize such models
 - what about other/more realistic structure?
- Tipping was inherently a *statistical* phenomenon
 - probabilistic nature of connectivity patterns
 - probabilistic nature of disease spread
 - model *likely* properties of a large *set* of possible outcomes
 - can model either inherent randomness or variability
- Formalizing tipping in the forest fire demo:
 - might let grid size $N \rightarrow$ infinity, look at fixed values of p
 - is there a threshold value q :
 - $p < q \rightarrow$ expected fraction burned $< 1/10$
 - $p > q \rightarrow$ expected fraction burned $> 9/10$

“Structural Virality” Goel, Anderson, Hofman, Watts

- Every video, news story, image, or petition posted to Twitter over 12 months (1.4 B observations)
 - Restrict to “popular” cascades (> 100 RTs; ~350K events)
- For each event, can quantify its “structural virality”
 - Average Pairwise Shortest Path Length
 - Ranges from
 - ≈ 2 (“broadcast”)
 - $\sim \log(N)$ (“viral”)
- For these “popular” events can ask:
 - What diversity do we see with respect to structure?
 - What is the relationship between size and structural virality?















“Broadcast”




“Viral”

Diversity of Structural Virality



Popular ≠ Viral



Popularity driven mostly by the size of the largest broadcast

